
Comet Prospects for 2018

Of the three periodic comets that should come into binocular range, two may be visible to the naked eye. The other is a Halley type comet last seen at its 1980 return, which is likely to be a well condensed object and therefore easier to see than some of the more diffuse periodic comets.

This draft version was updated on 2017 January 1 up to 2016 X1.

These predictions focus on comets that are likely to be within range of visual observers, though comets often do not behave as expected and can spring surprises. Members are encouraged to make visual magnitude estimates, particularly of periodic comets, as long term monitoring over many returns helps understand their evolution. Please submit your magnitude estimates in ICQ format. Guidance on visual observation and how to submit estimates is given in the BAA Observing Guide to Comets. Drawings are also useful, as the human eye can sometimes discern features that initially elude electronic devices.

Theories on the structure of comets suggest that any comet could fragment at any time, so it is worth keeping an eye on some of the fainter comets, which are often ignored. They would make useful targets for those making electronic observations, especially those with time on instruments such as the Faulkes telescopes. Such observers are encouraged to report electronic visual equivalent magnitude estimates via COBS. When possible use a waveband approximating to Visual or V magnitudes. These estimates can be used to extend the visual light curves, and hence derive more accurate absolute magnitudes. Such observations of periodic comets are particularly valuable as observations over many returns allow investigation into the evolution of comets.

In addition to the information in the BAA Handbook and on the Section web pages, ephemerides for the brighter observable comets are published in the *Circulars*, and ephemerides for new and currently observable comets are on the JPL, CBAT and Seiichi Yoshida's web pages. The BAA Observing Guide to Comets is available from the BAA Office.

21P/Giacobini-Zinner is the parent comet of the October Draconid meteors. On this return no significant activity from dust trail encounters is expected. The comet was first discovered by Michael Giacobini at Nice observatory in December 1900 and was thought to have a period of 6.8 years. The next two returns were expected to be difficult to observe, but in October 1913, Ernst Zinner, of Bamberg, Germany, discovered a comet whilst observing variable stars in Scutum. This turned out to be the same comet, but the period had been incorrectly determined and was actually 6.5 years. The comet was missed at three unfavourable returns, so the 2012 return was the fifteenth apparition of the comet. This is a very good apparition and at its closest in September the comet will be 0.39 au from the Earth. It will come within visual range in June and brightens rapidly. It could be a naked eye object by the end of August and remain visible to the naked eye throughout September. By then it is moving south and back into the morning sky, but keen UK observers should be able to follow it until the end of October. It passes relatively near quite a few open clusters. First, it is some 4° from the cluster M39 on July 4, but is much fainter than it. It passes just over a degree from open cluster

NGC 1502 around August 27, then M37 around September 13 and M35 a week later. NGC 2264 comes on September 26 and NGC 2301 at the beginning of October, with M50 and NGC 2343 a few days later and NGC 2362 at the end of the month.

29P/Schwassmann-Wachmann is an annual comet that has outbursts, which over the last decade seem to have become more frequent, though this could just reflect more intense coverage. Richard Miles has developed a theory that suggests that these outbursts are in fact periodic, and arise from at least four independent active areas on the slowly rotating nucleus. The activity of the active areas evolves with time. The comet is an ideal target for electronic observations and it should be observed at every opportunity. It is in solar conjunction in February. The comet creeps into the northern celestial sphere around the time of opposition in September when it is in Pisces.

Halley type comet **38P/Stephan-Oterma** was observed at its last return by the Section in 1980/81, when it reached 9th magnitude. Sky conditions were very different then when I was able to observe the well condensed comet in 10x80B from Cambridge Observatory. The comet was actually discovered by Jerome Coggia at the Marseilles Observatory, but the credit was taken by the Observatory Director, E J M Stephan, who obtained the first accurate position. It was then lost until a comet found in 1942 by Liisi Oterma at Turku, Finland was computed to be a return of Stephan's comet. It should come within visual range in September as it brightens rapidly on its way to the November perihelion. It will be at its brightest in November, when it is in the late evening sky and remains well placed into 2019.

Carl A Wirtanen discovered **46P/Wirtanen** at Lick in 1948. It is in a chaotic orbit, and its perihelion distance was much reduced due to approaches to Jupiter in 1972 and 84. It has been reported to outburst, but BAA data suggests that it was just rejuvenated after the perihelion distance was reduced. A December perihelion gives a close approach to the Earth and this will be achieved this year, when the comet passes 0.078 au from us. The comet may come into visual range in August, but it is then at a southern declination. UK visual observers may have a brief observing window in the early hours in September, when although it is still moving south it has brightened sufficiently that it may be visible above the sky glow. Otherwise we have to wait until the comet begins its rush north in November, when it may already be within binocular range. It is conveniently placed in our evening sky throughout December, when it may be visible to the naked eye. Indeed, the coma is likely to be very large due to its closeness, and the naked eye will be the best detector to use! The comet remains well placed as it fades into 2019 March.

48P/Johnson was discovered by Ernest Johnson at the Union Observatory in South Africa in 1949, following a very close approach to Jupiter in 1931. It is now in a stable orbit between Mars and Jupiter and no close approaches are predicted for some centuries. At favourable apparitions, such as its first two returns, it has reached 13th magnitude. The next three returns were unfavourable, with the comet reported to reach only 18th magnitude. Returns have now improved, and this is the closest perihelion passage for over a century. At the last return it reached 14th magnitude and it could be at least a couple of magnitudes brighter this time, though it will be best seen from the Southern Hemisphere.

185P/Petriew makes its fourth return, and although predicted to reach 11th magnitude, it will be a difficult object low down in the evening sky in the first few months of the year.

P/PANSTARRS (2013 CU₁₂₉) is an intrinsically faint comet, however it makes a relatively close pass of 0.24 au to the Earth in July when it may reach 11th magnitude. It is at a southern declination at this time, so will be a southern hemisphere target.

2016 M1 (PanSTARRS) is in solar conjunction in January and is heading towards a high southern declination as it emerges. For southern observers it should be picked up in early March and will be at its brightest of around 9th magnitude throughout April and May.

2016 R2 (PanSTARRS) could be 9th magnitude at the start of the year, however the magnitude parameters are currently uncertain and it might be brighter. It is well placed for observing by northern hemisphere observers. Despite not being at perihelion until May it will slowly fade as its increasing distance from the Earth compensates for its decreasing distance from the Sun. It might be 12th magnitude by the end of the year.

The other periodic and parabolic comets that are at perihelion during 2018 are unlikely to become brighter than 11th magnitude or are poorly placed. Ephemerides for these can be found on the CBAT or other WWW pages. Several D/ comets have predictions for a return, though searches at favourable returns in the intervening period have failed to reveal the comets and the orbits will have been perturbed by Jupiter. There is however always a chance that they will be rediscovered accidentally by one of the Sky Survey patrols.

Looking ahead to 2019 (or strictly 2020), there is only one additional periodic comet that is likely to be of significance. This is comet 289P/Blanpain, which was observed in 1819, then lost until re-discovered by the Catalina Sky Survey in 2003, when it passed 0.025 au from the Earth. It makes another close pass at 0.089 au in early 2020, when it might reach 5th magnitude and is well placed throughout the period when it is close to us. Two of the comets described above will also be on view.

Comets reaching perihelion in 2018

Comet	T	q	P	N	H ₁	K ₁	Peak mag
D/Brooks (1886 K1)	Oct 11.8	1.88	6.69	1			
D/Denning (1894 F1)	Feb 4.0	1.66	9.81	1			
D/Swift (1895 Q1)	Jun 28.0	1.39	7.18	1			
P/LINEAR-Skiff (2001 R6)	Oct 4.2	2.19	8.55	1	13.0	10.0	17
P/Catalina (2011 CR ₄)	Jun 22.6	2.52	6.58	1	13.0	5.0	16
P/PANSTARRS (2013 CU ₁₂₉)	Jun 24.3	0.80	4.88	1	15.2	10.0	11
P/Catalina-PANSTARRS (2013 R3)	Dec 6.4	2.20	5.28	1	14.0	10.0	18
PANSTARRS (2015 O1)	Feb 19.1	3.73			6.5	10.0	15
3D/Biela	Feb 23.3	0.80	6.57	6			
3D/Biela-A	Oct 3.6	0.81	6.60	2			
21P/Giacobini-Zinner	Sep 10.3	1.01	6.54	15	5.4	30.8	4
26P/Grigg-Skjellerup	Oct 1.8	1.08	5.23	20	12.0	40.0	15
37P/Forbes	May 4.1	1.61	6.43	11	9.5	10.0	12
38P/Stephan-Oterma	Nov 11.0	1.59	38.0	3	3.5	30.0	9
46P/Wirtanen	Dec 13.0	1.06	5.44	11	8.2	15.9	3

48P/Johnson	Aug 12.2	2.00	6.54	10	5.6	15.0	10
49P/Arend-Rigaux	Jul 15.5	1.43	6.74	10	9.6	10.0	13
59P/Kearns-Kwee	Sep 16.8	2.36	9.53	6	7.5	20.0	16
60P/Tsuchinshan	Dec 11.2	1.62	6.58	8	10.5	15.0	14
64P/Swift-Gehrels	Jun 14.4	1.38	9.34	6	9.0	20.0	13
66P/du Toit	May 19.1	1.29	14.9	3	12.0	9.5	13
74P/Smirnova-Chernykh	Jan 26.7	3.54	8.48	6	5.0	15.0	15
79P/du Toit-Hartley	Sep 13.3	1.12	5.05	6	14.0	15.0	16
82P/Gehrels	Jun 28.6	3.63	8.43	5	7.5	15.0	18
105P/Singer Brewster	Aug 10.3	2.04	6.46	5	12.5	15.0	18
107P/Wilson-Harrington	May 23.5	0.97	4.26	10	15.0	5.0	16
125P/Spacewatch	Aug 28.0	1.52	5.52	5	15.5	10.0	19
130P/McNaught-Hughes	Jan 21.8	1.82	6.22	4	12.5	10.0	17
133P/Elst-Pizarro	Sep 20.8	2.66	5.62	5	12.0	10.0	17
137P/Shoemaker-Levy	Dec 13.4	1.93	9.61	3	14.5	10.0	18
143P/Kowal-Mrkos	May 7.3	2.53	8.91	3	14.0	5.0	17
159P/LONEOS	May 22.8	3.63	14.2	2	10.0	10.0	18
164P/Christensen	May 31.4	1.69	7.01	3	11.0	10.0	15
169P/NEAT	Apr 29.6	0.60	4.20	6	16.0	5.0	16
185P/Petrew	Jan 27.7	0.93	5.46	3	10.7	19.6	11
187P/LINEAR	May 26.6	3.88	9.86	2	9.0	10.0	17
197P/LINEAR	Jan 28.8	1.06	4.85	3	16.5	5.0	18
198P/ODAS	Dec 13.9	2.01	6.84	3	10.5	15.0	15
235P/LINEAR	Mar 17.5	2.73	7.97	2	12.0	10.0	18
240P/NEAT	May 16.2	2.13	7.62	2	12.0	10.0	18
243P/NEAT	Aug 26.0	2.45	7.50	2	12.5	10.0	17
245P/WISE	Feb 8.0	2.16	8.06	2	14.0	10.0	19
247P/LINEAR	Dec 2.1	1.49	7.91	2	17.5	5.0	18
250P/Larson	Feb 1.5	2.21	7.21	3	14.5	10.0	18
253P/PANSTARRS	May 7.7	2.04	6.46	3	14.5	10.0	20
267P/LONEOS	Jul 22.3	1.24	5.76	2	19.5	10.0	20
282P/PANSTARRS	Apr 11.7	2.40	5.59	1	15.0	10.0	20
300P/Catalina	Nov 2.1	0.83	4.44	3	17.5	10.0	17
Lemmon (2015 XY ₁)	Apr 29.9	7.93			5.5	10.0	19
PanSTARRS (2016 M1)	Aug 9.8	2.21			5.0	10.0	9
PanSTARRS (2016 N6)	Jul 18.3	2.67			7.0	10.0	14
Kowalski (2016 Q4)	Jan 29.1	7.08			7.0	10.0	19
PanSTARRS (2016 R2)	May 16.7	2.64			3.0	10.0	9

The date of perihelion (T), perihelion distance (q), period (P), the number of previously observed returns (N), the magnitude parameters H_1 and K_1 and the brightest magnitude (which must be regarded as uncertain) are given for each comet. 3D has not been seen since 1852. The magnitudes, orbits, and in particular the time of perihelion of the single apparition D/ comets are uncertain.

Note: $m_1 = H_1 + 5.0 * \log(d) + K_1 * \log(r)$

References and sources

Belyaev, N. A., Kresak, L., Pittich, E. M. and Pushkarev, A. N., *Catalogue of short Period Comets*, Bratislava (1986).

Comet Observations Database (COBS) <http://www.cobs.si/> (Accessed 2016 October)

Comet Orbit Home Page (Kazuo Kinoshita) at <http://jcometobs.web.fc2.com/index.html> (Accessed 2015 October)

Jenniskens, P. *Meteor Showers and their Parent Comets*. Cambridge University Press (2006).

JPL Small-Body Database Browser <http://ssd.jpl.nasa.gov/sbdb.cgi#top> (Accessed 2016 October)

Kozlov, E. A., Medvedev, Y. D., Pittichova, J., and Pittich, E. M. *Catalogue of short Period Comets, 2nd edition*, (<http://astro.savba.sk/cat/>) (2003).

Kronk, G. W., *Cometographia*, Cambridge University Press, (1999, 2004, 2007, 2009, 2010) and <http://www.cometography.com> (Accessed 2016 October).

Marsden, B. G. and Williams, G. V. *Catalogue of Cometary Orbits*, 17th edition, IAU MPC/CBAT, (2008).

Minor Planet Electronic Circulars

Nakano Notes at <http://www.oaa.gr.jp/~oaacs/nk/> (Accessed 2016 October)

Shanklin, J. D., *Observing Guide to Comets*, 3rd edition (2013)

Jonathan Shanklin