

Stellar Dynamics and Structure of Galaxies

Derivation of potential from density distribution

Martin Haehnelt^{*}
haehnelt@ast.cam.ac.uk

Institute of Astronomy

Michaelmas Term 2018

^{*} based on slides prepared by Vasily Belokurov and lecture notes by Jim Pringle

Outline I

① Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

② Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow spherical density

Potentials from density distribution

Poisson's Equation

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Poisson's equation relates $\rho(\mathbf{r})$ to $\Phi(\mathbf{r})$.

Also covered in the Astrophysical Fluid Dynamics course

To determine the force due to a given density distribution $\rho(\mathbf{r}')$ we split it into many point masses of size

$$dm' = \rho(\mathbf{r}')d^3\mathbf{r}' \quad \text{at } \mathbf{r}'$$

Newtonian gravity is linear, so just add up the forces

$$\mathbf{f}(\mathbf{r}) = - \int \frac{Gdm'}{|\mathbf{r} - \mathbf{r}'|^3}(\mathbf{r} - \mathbf{r}')$$

or since we want the total potential add up the individual contributions

$$\Phi(\mathbf{r}) = \int \int \int \frac{G\rho(\mathbf{r}')d^3\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|}$$

As an exercise, show that $\nabla_r \frac{1}{|\mathbf{r}' - \mathbf{r}|} = \frac{\mathbf{r}' - \mathbf{r}}{|\mathbf{r}' - \mathbf{r}|^3}$, and hence $\mathbf{f}(\mathbf{r}) = -\nabla\Phi$

Potentials from density distribution

Poisson's Equation

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Consider

$$\nabla^2 \Phi(\mathbf{r}) = - \int \int \int G \rho(\mathbf{r}') \nabla_{\mathbf{r}}^2 \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} \right) d^3 \mathbf{r}'$$

\Rightarrow need $\nabla_{\mathbf{r}}^2 \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} \right)$.

To keep the algebra simple move the origin to \mathbf{r}' (and move back later)

for those who want everything in full generality, see Binney & Tremaine

So we need $\nabla^2 \left(\frac{1}{r} \right)$. For $r \neq 0$,

$$\nabla^2 \left(\frac{1}{r} \right) = \frac{1}{r^2} \frac{d}{dr} \left[r^2 \frac{d}{dr} \left(\frac{1}{r} \right) \right] = 0 \quad \text{trivially}$$

Potentials from density distribution

Poisson's Equation

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

But at $r = 0$ $\nabla^2\left(\frac{1}{r}\right)$ is undefined.

You've seen that sort of thing before. Recall that the Dirac δ -function $\delta(x)$ satisfies $\int_{-\epsilon}^{\epsilon} \delta(x) dx = 1$

So now ask: what is the volume integral of $\nabla^2\left(\frac{1}{r}\right)$ over a small volume V containing the origin?

$$\begin{aligned} \int \int \int_V \nabla^2 \left(\frac{1}{r} \right) d^3V &= \int \int \int_V \nabla \cdot \left[\nabla \left(\frac{1}{r} \right) \right] d^3V \text{ by definition} \\ &= \int \int_S \hat{\mathbf{n}} \cdot \left[\nabla \left(\frac{1}{r} \right) \right] d^2S \end{aligned} \quad (2.1)$$

Divergence theorem ($\hat{\mathbf{n}}$ - outward normal) $\int_V d^3\mathbf{x} \nabla \cdot \mathbf{F} = \int_S \hat{\mathbf{n}} \cdot \mathbf{F}$

Potentials from density distribution

Poisson's Equation

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Take V to be a sphere, so $\hat{\mathbf{n}} = \hat{\mathbf{r}}$, $d^2S = r^2 \sin \theta d\theta d\phi$, and have $\nabla(1/r) = -\frac{1}{r^2}\hat{\mathbf{r}}$. Then

$$\begin{aligned} \int \int \int_V \nabla^2 \left(\frac{1}{r} \right) d^3V &= - \int_0^{2\pi} d\phi \int_0^\pi \sin \theta d\theta \\ &= -4\pi \end{aligned} \quad (2.2)$$

Since the integral is -4π , and is non-zero only at $r = 0$, we must therefore have

$$\nabla^2 \left(\frac{1}{r} \right) = -4\pi\delta(\mathbf{r})$$

or, going back to the general origin,

$$\nabla^2 \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} \right) = -4\pi\delta(\mathbf{r} - \mathbf{r}')$$

Potentials from density distribution

Poisson's Equation

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Hence

$$\begin{aligned}\nabla^2\Phi(\mathbf{r}) &= -G \int \int \int \rho(\mathbf{r}') \nabla^2 \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} \right) d^3\mathbf{r}' \\ &= 4\pi G \int \int \int \rho(\mathbf{r}') \delta(\mathbf{r} - \mathbf{r}') d^3\mathbf{r}' \\ &= 4\pi G \rho(\mathbf{r})\end{aligned}\tag{2.3}$$

$$\nabla^2\Phi(\mathbf{r}) = 4\pi G \rho(\mathbf{r})$$

Poisson's Equation

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Gauss's Theorem

Application of the Divergence Theorem to the Poisson's Equation

“The integral of the normal component of $\nabla\Phi$ over any closed surface equals $4\pi G$ times the mass enclosed within that surface”

To prove this simply take Poisson's equation and integrate over a volume V containing a mass M .

$$\begin{aligned}
 4\pi G \int \rho d^3\mathbf{r} = 4\pi GM &= \int \nabla^2\Phi d^3\mathbf{r} \\
 &= \int \nabla \cdot \nabla\Phi d^3\mathbf{r} \\
 &= \int \nabla\Phi \cdot \hat{\mathbf{n}} d^2S \quad (2.4)
 \end{aligned}$$

where the last step follows from the divergence theorem.

Edwin Hubble's classification of galaxies

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

EXTRA-GALACTIC NEBULAE¹

By EDWIN HUBBLE

ABSTRACT

This contribution gives the results of a statistical investigation of 400 extra-galactic nebulae for which Holtschek has determined total visual magnitudes. The list is complete for the brighter nebulae in the northern sky and is representative to 12.5 mag. or fainter.

The classification employed is based on the forms of the photographic images. About 3 per cent are irregular, but the remaining nebulae fall into a sequence of type forms characterized by rotational symmetry about dominating nuclei. The sequence is composed of two sections, the elliptical nebulae and the spirals, which merge into each other.

Luminosity relations.—The distribution of magnitudes appears to be uniform throughout the sequence. For each type or stage in the sequence, the total magnitudes are related to the logarithms of the maximum diameters by the formula,

$$m_T = C - 5 \log d,$$

Astrophysical Journal, 64, 321-369
(1926)

Edwin Hubble's classification of galaxies

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

II. Extra-galactic nebulae:

A. Regular:

1. Elliptical..... E_n
 ($n=1, 2, \dots, 7$ indicates the ellipticity
 of the image without the decimal point)

{	N.G.C. 3379	E_0
	221	E_2
	4621	E_5
	2117	E_7

2. Spirals:

	Symbol	Example
a) Normal spirals.....	S
(1) Early.....	Sa	N.G.C. 4594
(2) Intermediate.....	Sb	2841
(3) Late.....	Sc	5457
b) Barred spirals.....	SB
(1) Early.....	SBa	N.G.C. 2859
(2) Intermediate.....	SBb	3351
(3) Late.....	SBc	7479
B. Irregular.....	Irr	N.G.C. 4449

Edwin Hubble's classification of galaxies

Early Types

Potentials from density distribution


Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials



Fundamental plane exists that ties surface brightness, size and LOS velocity dispersion

Edwin Hubble's classification of galaxies

Spirals

Potentials from density distribution


Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials



Tully-Fisher law exists that ties together circular speed and luminosity

Edwin Hubble's classification of galaxies

Irregulars

Potentials from density distribution


Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials



Edwin Hubble's classification of galaxies

The Tuning Fork

Potentials from density distribution


Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials



Edwin Hubble's classification of galaxies

The Three Pioneers

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems


Profiles and potentials



Albert Einstein, Edwin Hubble, and Walter Adams in 1931 at the Mount Wilson Observatory 100" telescope, in the San Gabriel Mountains of southern California.

Edwin Hubble's classification of galaxies

Galaxy Luminosity Function



In any environment, dwarfs dominate!

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Deriving potentials of spherical systems

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's

classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

we can take $\rho(r) = \rho(r)$

In spherical polars

$$\nabla^2 \Phi(r) = \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d}{dr} \Phi \right) = \frac{1}{r} \frac{d^2}{dr^2} (r\Phi)$$

Exercise: show the last equality is true

So

$$\nabla^2 \Phi = 4\pi G\rho$$

becomes

$$\frac{1}{r} \frac{d^2}{dr^2} (r\Phi) = 4\pi G\rho,$$

and, given ρ we can solve for $\Phi(r)$.

Deriving potentials of spherical systems

Homogeneous Sphere

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's

classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

(a) Homogeneous sphere: $\rho(r) = \rho_0$ for $0 < r < r_0$, and $\rho(r) = 0$ for $r > r_0$.

So for $r < r_0$, have

$$\begin{aligned} \frac{1}{r} \frac{d^2}{dr^2} (r\phi) &= 4\pi G\rho_0 \\ \frac{d^2}{dr^2} (r\phi) &= 4\pi G\rho_0 r \\ \frac{d}{dr} (r\phi) &= 2\pi G\rho_0 r^2 + A \\ r\phi &= \frac{2}{3}\pi G\rho_0 r^3 + Ar + B \\ \phi(r) &= \frac{2}{3}\pi G\rho_0 r^2 + A + \frac{B}{r} \end{aligned} \quad (2.5)$$

Deriving potentials of spherical systems

Homogeneous Sphere

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's

classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

$$\Phi(r) = \frac{2}{3}\pi G\rho_0 r^2 + A + \frac{B}{r}$$

Require that Φ is finite at $r = 0$, else there is a point mass there, and so $B = 0$.

$$\Rightarrow \Phi(r) = \frac{2}{3}\pi G\rho_0 r^2 + A \text{ for } 0 < r < r_0.$$

For $r > r_0$ have

$$\frac{1}{r} \frac{d^2}{dr^2} (r\Phi) = 0$$

$$\Rightarrow r\Phi = Cr + D$$

$$\Phi(r) = C + \frac{D}{r}$$

WLOG¹ let $\Phi \rightarrow 0$ as $r \rightarrow \infty$ (this is just choosing the zero point of the potential).

$$\Rightarrow \Phi(r) = \frac{D}{r} \text{ for } r_0 < r$$

¹WLOG=Without Loss Of Generality

Deriving potentials of spherical systems

Homogeneous Sphere

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's

classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

$$\Phi(r) = \frac{2}{3}\pi G\rho_0 r^2 + A \text{ for } 0 < r < r_0$$

$$\Phi(r) = \frac{D}{r} \text{ for } r_0 < r$$

Also require Φ to be continuous at $r = r_0$, since $\nabla\Phi = \text{force}$ is finite there, and $\frac{d\Phi}{dr}$ also continuous (else $\nabla^2\Phi = 4\pi G\rho$ is infinite there).

$$\Rightarrow \frac{2}{3}\pi G\rho_0 r_0^2 + A = \frac{D}{r_0}$$

and

$$\frac{4}{3}\pi G\rho_0 r_0 = -\frac{D}{r_0^2}$$

 \Rightarrow

$$D = -\frac{4}{3}\pi G\rho_0 r_0^3$$

and

$$A = -2\pi G\rho_0 r_0^2$$

Deriving potentials of spherical systems

Homogeneous Sphere

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Hence

Potential of a homogeneous sphere

$$\begin{aligned}\Phi(r) &= \frac{2}{3}\pi G\rho_0(r^2 - 3r_0^2) \quad 0 < r < r_0 \\ &= -\frac{4}{3}\pi G\rho_0 r_0^3/r \quad r_0 < r\end{aligned}\quad (2.6)$$

Note: Outside the sphere $\Phi = -\frac{GM}{r}$ as expected, where $M = \frac{4}{3}\pi\rho_0 r_0^3$.

Newton's 2nd theorem: "Outside a closed spherical shell of matter, the gravitational potential is as if all the mass were at a point at the centre"

Deriving potentials of spherical systems

Spherical Shell

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's

classification of galaxies

Deriving potentials of

spherical systems

Profiles and potentials

(b) Spherical shell $\rho(r) = \rho_0$ for $r_1 < r < r_2$ and $\rho(r) = 0$ otherwise. Newtonian gravity is linear, so this is the same as

(1) a uniform sphere density ρ_0 , radius r_2

PLUS

(2) a uniform sphere density $-\rho_0$, radius r_1 .

So we can write the answer down. It is

$$\begin{aligned}
 \Phi(r) &= \frac{2}{3}\pi G\rho_0(r^2 - 3r_2^2) - \frac{2}{3}\pi G\rho_0(r^2 - 3r_1^2) \quad 0 < r < r_1 \\
 &= \frac{2}{3}\pi G\rho_0(r^2 - 3r_2^2) + \frac{4}{3}\pi G\rho_0 r_1^3/r \quad r_1 < r < r_2 \\
 &= -\frac{4}{3}\pi G\rho_0 r_2^3/r + \frac{4}{3}\pi G\rho_0 r_1^3/r \quad r_2 < r
 \end{aligned} \tag{2.7}$$

Deriving potentials of spherical systems

Spherical Shell

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's

classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Notes:

(1) Inside the cavity $0 < r < r_1$:

$$\Phi(r) = \frac{2}{3}\pi G\rho_0(r^2 - 3r_2^2) - \frac{2}{3}\pi G\rho_0(r^2 - 3r_1^2)$$

$\Phi = \text{constant}$ since the r^2 terms cancel. Therefore there is no force due to an external spherically symmetric mass distribution

Newton's first theorem

(2) Outside the shell $r > r_2$: $\Phi(r) = -\frac{4}{3}\pi G\rho_0 r_2^3/r + \frac{4}{3}\pi G\rho_0 r_1^3/r$

$$\Phi = -\frac{GM_{\text{shell}}}{r}$$

where $M_{\text{shell}} = \frac{4}{3}\pi\rho_0(r_2^3 - r_1^3)$ is the mass in the shell

Deriving potentials of spherical systems

Shells Galore

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's

classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Since Newtonian gravitational potentials add linearly, we can calculate the potential at r due to an arbitrary spherically symmetric $\rho(r)$ by adding contributions from shells inside and outside r .

Mass in shell of thickness dr' and radius r' is

$$4\pi r'^2 \rho(r') dr'$$

The potential inside a shell is constant, so we can evaluate it anywhere - easiest is just inside the shell, where

$$\Phi = - \frac{4\pi G r'^2 \rho(r') dr'}{r'}$$

(from $-GM/r$).

Deriving potentials of spherical systems

Shells Galore

Potentials from density distribution

Poisson's Equation

Gauss's Theorem

Edwin Hubble's

classification of galaxies

Deriving potentials of spherical systems

Profiles and potentials

Thus, at any r , we have:

$$\Phi(r) = -\frac{4\pi G}{r} \int_0^r r'^2 \rho(r') dr' - 4\pi G \int_r^\infty r' \rho(r') dr'$$

where the first term is from shells inside r , and the second from shells outside r (to get $\Phi(\infty) = 0$).

Potential of an arbitrary spherical distribution

$$\Phi(r) = -4\pi G \left[\frac{1}{r} \int_0^r r'^2 \rho(r') dr' + \int_r^\infty r' \rho(r') dr' \right] \quad (2.8)$$

Profiles and potentials

Modified Hubble profile

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

If a galaxy has a spherical luminosity density

$$j(r) = j_0 \left(1 + \left(\frac{r}{a} \right)^2 \right)^{-\frac{3}{2}} \quad (2.9)$$

then the surface brightness distribution is the projection of this on the plane of the sky

$$I(R) = 2 \int_0^{\infty} j(z) dz \quad (2.10)$$

Now $r^2 = R^2 + z^2$, so

$$I(R) = 2j_0 \int_0^{\infty} \left[1 + \left(\frac{R}{a} \right)^2 + \left(\frac{z}{a} \right)^2 \right]^{-\frac{3}{2}} dz \quad (2.11)$$

Profiles and potentials

Modified Hubble profile

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

Let $y = z/\sqrt{a^2 + R^2}$, and then

$$1 + \left(\frac{R}{a}\right)^2 + \left(\frac{z}{a}\right)^2 = \frac{1}{a^2} (a^2 + R^2 + z^2) = \frac{(a^2 + R^2)}{a^2} (1 + y^2) \quad (2.12)$$

$$\Rightarrow I(R) = 2j_0 \left(\frac{a^2}{a^2 + R^2}\right)^{\frac{3}{2}} \int_0^\infty \frac{\sqrt{a^2 + R^2} dy}{(1 + y^2)^{\frac{3}{2}}} \quad (2.13)$$

$$= 2j_0 \frac{a^3}{a^2 + R^2} \int_0^\infty \frac{dy}{(1 + y^2)^{\frac{3}{2}}} \quad (2.14)$$

Can be evaluated by setting $y = \tan x$, so $dy = \sec^2 x dx$, and the integral becomes

$$\int_0^{\frac{\pi}{2}} \frac{\sec^2 x dx}{(\sec^2 x)^{\frac{3}{2}}} = \int_0^{\frac{\pi}{2}} \cos x dx = \sin \frac{\pi}{2} - \sin 0 = 1 \quad (2.15)$$

Profiles and potentials

Modified Hubble profile

and hence

$$\begin{aligned}
 I(R) = 2 \int_0^\infty j(z) dz &= 2j_0 \int_0^\infty \left[1 + \left(\frac{R}{a}\right)^2 + \left(\frac{z}{a}\right)^2 \right]^{-\frac{3}{2}} dz \\
 &= 2j_0 \frac{a^3}{a^2 + R^2} \int_0^\infty \frac{dy}{(1 + y^2)^{\frac{3}{2}}} = \frac{2j_0 a}{1 + \left(\frac{R^2}{a^2}\right)} \quad (2.16)
 \end{aligned}$$

This profile is quite a good fit to elliptical galaxies - it is similar to the Hubble profile.

Now ask: assuming a fixed mass-to-light ratio Υ , what is the potential?

Assume

$$\rho(r) = \frac{\rho_0}{\left[1 + \left(\frac{r}{a}\right)^2 \right]^{\frac{3}{2}}} \quad (2.17)$$

where $\rho_0 = \Upsilon j_0$.

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

Profiles and potentials

Modified Hubble profile

Let's use Poisson's equation $\nabla^2 \Phi = 4\pi G \rho \Rightarrow \frac{d^2}{dr^2} r\Phi = 4\pi G r \rho$

$$\begin{aligned} \frac{1}{4\pi G} \frac{d^2}{dr^2} (r\Phi) &= \frac{\rho_0 r}{\left(1 + \frac{r^2}{a^2}\right)^{\frac{3}{2}}} \\ \frac{1}{4\pi G} \frac{d}{dr} (r\Phi) &= \rho_0 \int \frac{r dr}{\left(1 + \frac{r^2}{a^2}\right)^{\frac{3}{2}}} \\ &= \frac{\rho_0 a^2}{2} \int \frac{2r dr/a^2}{\left(1 + r^2/a^2\right)^{\frac{3}{2}}} \end{aligned} \quad (2.18)$$

Let $u = 1 + r^2/a^2$, then $du = \frac{2r}{a^2} dr$

Profiles and potentials

Modified Hubble profile

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

$$u = 1 + r^2/a^2$$

$$\rho(r) = \frac{\rho_0}{\left[1 + \left(\frac{r}{a}\right)^2\right]^{\frac{3}{2}}}$$

$$\frac{d^2}{dr^2} r\Phi = 4\pi G r \rho$$

And so

$$\begin{aligned} \frac{1}{4\pi G} \frac{d}{dr} (r\Phi) &= \frac{\rho_0 a^2}{2} \int \frac{du}{u^{\frac{3}{2}}} \\ &= -2 \frac{\rho_0 a^2}{2} \left(1 + \frac{r^2}{a^2}\right)^{-\frac{1}{2}} + A \end{aligned} \quad (2.19)$$

Then

$$\frac{r\Phi}{4\pi G} = Ar - \rho_0 a^3 \int \frac{dr}{\sqrt{a^2 + r^2}} \quad (2.20)$$

Then we have the fairly standard integral

$$\int \frac{dx}{\sqrt{a^2 + x^2}} = \ln(2\sqrt{a^2 + x^2} + 2x) \text{ or } \sinh^{-1} \left(\frac{x}{a}\right) \quad (2.21)$$

Profiles and potentials

Modified Hubble profile

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

So

$$\frac{r\Phi}{4\pi G} = Ar + B - \rho_0 a^3 \ln(2\sqrt{a^2 + r^2} + 2r) \quad (2.22)$$

$B = \rho_0 a^3 \ln(2a)$ as otherwise $1/r \rightarrow \infty$ as $r \rightarrow 0$ [i.e. no point mass at origin].

$$\Phi = 4\pi GA - 4\pi G\rho_0 a^3 \frac{\ln(2\sqrt{a^2 + r^2} + 2r) - \ln(2a)}{r} \quad (2.23)$$

Note that we can choose $A = 0$, and then $\Phi \rightarrow 0$ as $r \rightarrow \infty$ (but more slowly than $\frac{1}{r}$ due to infinite total mass).

The total mass within r is

$$M(r) = \int_0^r \frac{4\pi\rho_0 r^2 dr}{\left(1 + \frac{r^2}{a^2}\right)^{\frac{3}{2}}} \quad (2.24)$$

This is $\propto \ln r$ for large r , so diverges as $r \rightarrow \infty$.

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

Profiles and potentials

Power law density profile

$$\rho(r) = \rho_0 \left(\frac{a}{r}\right)^\alpha \quad (2.25)$$

$$\frac{d^2}{dr^2}(r\Phi) = 4\pi G\rho_0 a^\alpha r^{1-\alpha} \quad (2.26)$$

so

$$\frac{d}{dr}(r\Phi) = 4\pi G\rho_0 a^\alpha \frac{r^{2-\alpha}}{2-\alpha} + A \quad (2.27)$$

Profiles and potentials

Power law density profile

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

$$r\Phi = 4\pi G\rho_0 a^\alpha \frac{r^{3-\alpha}}{(2-\alpha)(3-\alpha)} + Ar + B \quad (2.28)$$

or

$$\Phi = -\frac{4\pi G\rho_0 a^\alpha r^{2-\alpha}}{(3-\alpha)(\alpha-2)} + A + \frac{B}{r} \quad (2.29)$$

$A = 0$ by setting zero, and $B = 0$ because no point mass at centre as usual.

Profiles and potentials

Power law density profile

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

$$\Phi = -\frac{4\pi G\rho_0 a^\alpha r^{2-\alpha}}{(3-\alpha)(\alpha-2)}$$

$$\rho(r) = \rho_0 \left(\frac{a}{r}\right)^\alpha$$

Notes:

(1) $\alpha < 3$ to get $M(r)$ finite at the origin (determine $\int 4\pi G\rho r^2 dr$ near origin).

(2) $\Phi \rightarrow 0$ at ∞ if $\alpha > 2$,

$$\Rightarrow 2 < \alpha < 3$$

$\alpha = 2$ gives spiral rotation curves (flat), from

$$v_c^2/r = \frac{d\Phi}{dr} (= -f_r) \Rightarrow v_c^2 \propto r^{2-\alpha}.$$

[Circular motion $\Rightarrow \ddot{r} \& \dot{r} = 0$, so $\ddot{r} - r\dot{\phi}^2 = -\frac{d\Phi}{dr}$ becomes, with

$$v_c = r\dot{\phi}, \frac{v_c^2}{r} = -\frac{d\Phi}{dr}. \text{ Then substituting } \Phi \text{ from equation (2.29) gives } v_c^2 \propto r^{2-\alpha}.]$$

$\alpha = 3$ gives elliptical galaxy profiles (mod. Hubble profile)

but all these models have infinite mass, since $M(r)$ diverges at large r

Projected density \rightarrow spherical density

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density


What we have done so far is to guess a luminosity density $j(r)$ (which we assume is proportional to the matter density $\rho(r)$) and formed the projected surface brightness $I(R)$ using the relation

$$I(R) = 2 \int_R^\infty \frac{j(r)rdr}{\sqrt{r^2 - R^2}} \quad (2.30)$$

and then check that $I(R)$ is a reasonable approximation to what is seen for our guessed density distribution.

$$r^2 = x^2 + R^2$$

$$dx = \frac{rdr}{\sqrt{r^2 - R^2}}$$



Projected density \rightarrow spherical density

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

OK, so

$$I(R) = 2 \int_R^\infty \frac{j(r)rdr}{\sqrt{r^2 - R^2}}$$

In fact, if $I(R)$ is known, then the equation above may be inverted to yield $j(r)$ directly, to yield

$$j(r) = -\frac{1}{2\pi r} \frac{d}{dr} \int_r^\infty \frac{I(R)RdR}{\sqrt{R^2 - r^2}}. \quad (2.31)$$

This is not quite pulled out of the air - it is a form of Abel's integral equation.

Projected density \rightarrow spherical density

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

We can simplify the form a bit if we set $t = R^2$ and $x = r^2$, and then we have

$$I(t) = \int_t^\infty \frac{j(x)dx}{(x-t)^{\frac{1}{2}}}$$

and then the inverse relation quoted becomes

$$j(y) = -\frac{1}{\pi} \frac{d}{dy} \int_y^\infty \frac{I(t)dt}{(t-y)^{\frac{1}{2}}}$$

If we look just at the RHS, and call it $h(y)$ for the moment, this is

$$h(y) = -\frac{1}{\pi} \frac{d}{dy} \int_y^\infty \frac{dt}{(t-y)^{\frac{1}{2}}} \int_t^\infty \frac{j(x)dx}{(x-t)^{\frac{1}{2}}}.$$

or

$$h(y) = -\frac{1}{\pi} \frac{d}{dy} \int_{t=y}^\infty \int_{x=t}^\infty \frac{dtj(x)dx}{(t-y)^{\frac{1}{2}}(x-t)^{\frac{1}{2}}}$$

Projected density \rightarrow spherical density

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

We now switch the order of the integration, remembering when doing so to change the limits of the integration so that we are integrating over the same area in the (x, t) -plane.

$$h(y) = -\frac{1}{\pi} \frac{d}{dy} \int_y^\infty j(x) dx \int_y^x \frac{dt}{(t-y)^{\frac{1}{2}}(x-t)^{\frac{1}{2}}}$$

The integral

$$\int_y^x \frac{dt}{(t-y)^{\frac{1}{2}}(x-t)^{\frac{1}{2}}} = \pi$$

and so what we called $h(y)$ is then seen to be equal to $j(y)$. So the result follows.

Projected density \rightarrow spherical density

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

[The statement that

$$S \equiv \int_y^x \frac{dt}{(t-y)^{\frac{1}{2}}(x-t)^{\frac{1}{2}}} = \pi$$

needs a bit more justification, or you can take it on trust.... For those who don't, we first change variables so the lower limit is zero, so $z = t - y$, and then

$$S = \int_0^{x-y} \frac{dz}{(x-y-z)^{\frac{1}{2}}z^{\frac{1}{2}}}$$

This invites yet another change of variables so that the upper limit is 1, i.e. $\zeta = \frac{z}{x-y} \Rightarrow z = (x-y)\zeta \Rightarrow x-y-z = (x-y)(1-\zeta) \Rightarrow$

$$S = \int_0^1 \frac{(x-y)d\zeta}{(x-y)^{\frac{1}{2}}(1-\zeta)^{\frac{1}{2}}(x-y)^{\frac{1}{2}}\zeta^{\frac{1}{2}}}$$

Projected density \rightarrow spherical density

So

$$\begin{aligned}
 S &= \int_0^1 \frac{d\zeta}{(1-\zeta)^{\frac{1}{2}} \zeta^{\frac{1}{2}}} & (2.32) \\
 &= \int_0^1 \frac{d\zeta}{(\zeta - \zeta^2)^{\frac{1}{2}}} \\
 &= \int_0^1 \frac{d\zeta}{\left(\frac{1}{4} - \left(\zeta - \frac{1}{2}\right)^2\right)^{\frac{1}{2}}} \\
 &= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{du}{\sqrt{\frac{1}{4} - u^2}} \\
 &= \int_{-1}^1 \frac{\frac{1}{2} dv}{\sqrt{\frac{1}{4} - \frac{v^2}{4}}} \\
 &= \int_{-1}^1 \frac{dv}{\sqrt{1 - v^2}}
 \end{aligned}$$

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

Projected density \rightarrow spherical density

Potentials from density distribution

Profiles and potentials

Modified Hubble profile

Power law density profile

Projected density \rightarrow
spherical density

Then since we know

$$\frac{d}{d\xi} \arcsin \xi = \frac{1}{\sqrt{1 - \xi^2}}$$

we have

$$\int_{-1}^1 \frac{dv}{\sqrt{1 - v^2}} = \arcsin v \Big|_{-1}^1 = \frac{\pi}{2} + \frac{\pi}{2} = \pi$$

]