

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Stellar Dynamics and Structure of Galaxies

Potentials due to axisymmetric distributions of matter

Martin Haehnelt^{*}
haehnelt@ast.cam.ac.uk

Institute of Astronomy

Michaelmas Term 2018

^{*} based on slides prepared by Vasily Belokurov and lecture notes by Jim Pringle

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

- ① Potentials due to axisymmetric matter distributions
 - Legendré Polynomials
 - Example
 - Motion of the Moon
- ② Potential due to thin disk
 - Bessel Functions
 - Poisson's equation
 - Summary of derivation of Φ for thin axisymmetric disk
 - Examples
 - Mestel disk
 - Exponential Disk
- ③ The Oort Constants: the rotation of our Galaxy
- ④ The Oort Constants and epicycles
- ⑤ The Rotation Curve of our Galaxy
- ⑥ Spiral Galaxy Rotation Curves

Potential from axisymmetric matter distribution

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles


The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Outside matter, $\rho = 0$, so $\nabla^2\Phi = 0$.

Axisymmetric, so independent of spherical polar ϕ ,

$$\Phi = \Phi(r, \theta) \text{ only}$$



Potential from axisymmetric matter distribution

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Outside matter, $\rho = 0$, so $\nabla^2\Phi = 0$.

Axisymmetric, so independent of spherical polar ϕ ,

$$\Phi = \Phi(r, \theta) \text{ only}$$

Then $\nabla^2\Phi = 0$ becomes

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \Phi}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \Phi}{\partial \theta} \right) = 0.$$

If we **assume** the function $\Phi(r, \theta)$ is separable, ie

$$\Phi(r, \theta) = \mathcal{R}(r) \times \Theta(\theta)$$

then, as substituting and dividing by $\mathcal{R}\Theta$,

$$\frac{1}{\mathcal{R}} \frac{d}{dr} \left(r^2 \frac{d\mathcal{R}}{dr} \right) + \frac{1}{\Theta \sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) = 0$$

i.e. **each term has to be constant (and one = - the other)**

Potential from axisymmetric matter distribution

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Let

$$\frac{1}{\mathcal{R}} \frac{d}{dr} \left(r^2 \frac{d\mathcal{R}}{dr} \right) = n(n+1) \quad (\text{say}) \quad (4.1)$$

$$\text{then } \frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) = -n(n+1)\Theta(\theta) \quad (4.2)$$

Set $\mu = \cos \theta$ ($-1 \leq \mu \leq 1$)

$$\frac{d}{d\mu} \left\{ (1 - \mu^2) \frac{d\Theta}{d\mu} \right\} + n(n+1)\Theta = 0 \quad (4.3)$$

since $d\mu = -\sin \theta d\theta$

– Legendres equation.

Want $\Theta(\theta)$ to be finite at $\theta = 0, \pi$ ($\mu = \pm 1$). And the only way this can happen is if $n = 0, 1, 2, \dots$ - a non-negative integer

Potential from axisymmetric matter distribution

Legendré Polynomials

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

$$\frac{d}{d\mu} \left\{ (1 - \mu^2) \frac{d\Theta}{d\mu} \right\} + n(n+1)\Theta = 0 \quad 4.3$$

Solution of (4.1) is

$$\mathcal{R}(r) = Ar^n + \frac{B}{r^{n+1}}$$

And (4.3) is the defining differential equation for Legendré polynomials, so we have

$$\text{and } \Theta(\theta) = CP_n(\cos \theta) \text{ for a given } n$$

Presumably you've seen Legendré polynomials before:

$$P_0(x) = 1$$


$$P_n(1) = 1 \forall n$$

$$P_1(x) = x$$

$$P_2(x) = \frac{1}{2} (3x^2 - 1)$$

Potential from axisymmetric matter distribution

Legendré Polynomials



Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential from axisymmetric matter distribution

Legendré Polynomials

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Recurrence relation:

$$(n + 1)P_{n+1}(x) = (2n + 1)xP_n(x) - nP_{n-1}(x) .$$

Orthogonal polynomials on $[-1, 1]$ i.e.

$$\int_{-1}^1 P_n(x)P_m(x) dx = 0 \quad \text{if } n \neq m$$

$$\left(= \frac{2}{2n + 1} \quad \text{if } n = m \right)$$

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n .$$

$P_n(x)$ is oscillatory: $P_n(x)$ has n zeros in $[-1,1]$ (so higher $n \Rightarrow$ more oscillations)

\Rightarrow Any function on $[-1,1]$ can be reproduced as a sum of these (cf Fourier series).

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Legendré and Fourier



The only known portrait of Legendré, recently unearthed, is found in the 1820 book *Album de 73 portraits-charge aquarelés des membres de l'Institut*, a book of caricatures of seventy-three famous mathematicians by the French artist Julien-Leopold Boilly

Potential from axisymmetric matter distribution

Legendré Polynomials

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Note then for general solution outside axisymmetric body

$$\mathcal{R}(r) = \frac{B}{r^{n+1}}$$

(want Φ finite as $r \rightarrow \infty$).


And if the body is symmetric about $\theta = \pi/2$ plane, then only need even n , since these are symmetric about $\mu = 0$ and the odd n ones are antisymmetric.

$$\begin{aligned} \Phi(r, \theta) &= \sum_{n=0}^{\infty} \frac{B_n P_n(\cos \theta)}{r^{n+1}} = \sum_{k=0}^{\infty} \frac{B_{2k} P_{2k}(\cos \theta)}{r^{2k+1}} \\ &= -\frac{GM}{r} + \frac{J_2}{r^3} \frac{1}{2} (3 \cos^2 \theta - 1) + \frac{J_4}{r^5} \left(\frac{35}{8} \cos^4 \theta - \frac{15}{4} \cos^2 \theta + \frac{3}{8} \right) \dots \end{aligned}$$

If the body is nearly spherical, treat J terms as perturbations.

Potential from axisymmetric matter distribution

Example



Potential due to a ring of matter
Mass M , radius a (in full generality)

Axisymmetric \Rightarrow no ϕ dependence $\Rightarrow \nabla^2 \Phi = 0$ has solution

$$\Phi = \sum_{n=0}^{\infty} \left(A_n r^n + \frac{B_n}{r^{n+1}} \right) P_n(\cos \theta)$$

So if we need to work out Φ somewhere useful to get the coefficients A_n and B_n .

Trick: let's try the easiest place - on the z axis.

At height z on the z -axis

$$\Phi = -G \int \frac{\rho d^3 \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} = -\frac{GM}{(a^2 + z^2)^{\frac{1}{2}}}$$

\uparrow usual sum

Potentials due to axisymmetric matter distributions

Legendre Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential from axisymmetric matter distribution

Example

For small z ,

$$\begin{aligned}\Phi &= -\frac{GM}{a} \left(1 + \frac{z^2}{a^2}\right)^{-\frac{1}{2}} \\ &= -\frac{GM}{a} \left\{1 - \frac{z^2}{2a^2} + \frac{3z^4}{8a^4} - \dots\right\}\end{aligned}$$

But on axis also $\theta = 0, \cos\theta = 1, r = z$ so

$$\Phi(r) = \sum_{n=0}^{\infty} \left(A_n z^n + \frac{B_n}{z^{n+1}} \right) \text{ there}$$

Match terms, and so for small $r < a$

$$\Phi = -\frac{GM}{a} \left\{ 1 - \frac{r^2}{2a^2} P_2(\cos\theta) + \frac{3}{8a^4} r^4 P_4(\cos\theta) \dots \right\}$$

Potentials due to axisymmetric matter distributions

Legendre Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential from axisymmetric matter distribution

Example

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon


Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



$$\Phi = -\frac{GM}{a} \left\{ 1 - \frac{r^2}{2a^2} P_2(\cos \theta) + \frac{3}{8a^4} r^4 P_4(\cos \theta) \dots \right\}$$

$$P_0(\cos \theta) = 1$$

$$P_2(\cos \theta) = \frac{1}{2}(3 \cos^2 \theta - 1)$$

$$P_4(\cos \theta) = \frac{1}{8}(35 \cos^4 \theta - 30 \cos^2 \theta + 3)$$

So if you want the potential close to the plane:

$$\theta \sim \pi/2 \quad r \rightarrow R, \quad \text{and } r \cos \theta = z \quad \text{or } \cos \theta = \frac{z}{R}$$

$$\Rightarrow \Phi(R, Z) \simeq -\frac{GM}{a} \left[1 - \frac{1}{4a^2} \{3z^2 - R^2\} + \dots \right]$$

Potential from axisymmetric matter distribution

Motion of the Moon



Can apply this to the motion of the Moon due to the time average of the Sun (axisymmetric part of the potential)

$$\Phi = -\frac{GM_{\oplus}}{R} - \frac{GM_{\odot}}{a} \left\{ 1 - \frac{1}{4a^2} (3z^2 - R^2) \right\}$$

$$\begin{aligned} R = R_{\oplus M} &= 4 \times 10^8 \text{ m} & a = R_{\odot \oplus} &= 1.5 \times 10^{11} \text{ m} \\ M_{\oplus} &= 6 \times 10^{24} \text{ kg} & M_{\odot} &= 2 \times 10^{30} \text{ kg} \end{aligned}$$

Mass of moon is about 1/81th that of the Earth, so it is the 'test particle'.

Ratio of Earth-Moon and Moon-Sun extra part is

$$\frac{GM_{\oplus}}{R} / \frac{GM_{\odot} R^2}{a^3} = M_{\oplus} a^3 / M_{\odot} R^3 \sim 200$$

so can treat the Moon-Sun force as a perturbation.

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential from axisymmetric matter distribution

Eclipse cycle

$R_{\oplus} = 6360 \text{ km}$ – need accurate alignment.

Moon orbit inclination to ecliptic 5° , eccentricity 0.055.

Moon's period:

- Siderial (relative to fixed star) $= 27^{\text{d}}.322 = 2\pi/\Omega$
- Synodical (new moon - new moon) $= 29^{\text{d}}.530 = 2\pi/(\Omega - \Omega_{\oplus})$
- Anomalistic month (perigee to perigee) $= 27^{\text{d}}.554 = 2\pi/\Omega_r = 2\pi/K$
- Nodical month (node to node) $= 27^{\text{d}}.212 = 2\pi/\mathcal{V}$

To get a solar eclipse we must have

- Moon at “new moon” - once every synodical month
- Moon passing through a node - once every nodical month
- For the best eclipse, moon at perigee - once every anomalistic month

The “saros” or eclipse cycle comes about because

223 synodic months = 6585.32 days

242 nodical months = 6585.36 days

239 anomalistic months = 6585.54 days

... so every 6585 days, or 18 years get a similar pattern of eclipses.

Potentials due to axisymmetric matter distributions

Legendré Polynomials

Example

Motion of the Moon

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions
Poisson's equation
Summary of derivation of Φ for thin axisymmetric disk
Examples


The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk



Let's use cylindrical polar, (R, ϕ, z) , coordinates

Expect $\Phi \equiv \Phi(R, z)$ and $\Phi(R, z) = \Phi(R, -z)$ by symmetry.

Outside disk $\nabla^2 \Phi = 0$.

$$\Rightarrow \frac{1}{R} \frac{\partial}{\partial R} \left(R \frac{\partial \Phi}{\partial R} \right) + \frac{\partial^2 \Phi}{\partial z^2} = 0$$

We solve this by separation of variables, letting

$$\Phi(R, z) = J(R)Z(z)$$

Potential due to thin disk

Bessel Functions

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

$$\frac{1}{R} \frac{\partial}{\partial R} \left(R \frac{\partial \Phi}{\partial R} \right) + \frac{\partial^2 \Phi}{\partial z^2} = 0$$

$$\Phi(R, z) = J(R)Z(z)$$

$$\Rightarrow Z(z) \frac{1}{R} \frac{d}{dR} \left(R \frac{dJ(R)}{dR} \right) + J(R) \frac{d^2 Z(z)}{dz^2} = 0$$

$$\Rightarrow \underbrace{\frac{1}{JR} \frac{d}{dR} \left(R \frac{dJ}{dR} \right)}_{\text{function of } R} = - \underbrace{\frac{1}{Z} \frac{d^2 Z}{dz^2}}_{\text{function of } z} = -k^2, \text{ say}$$

$$\Rightarrow \frac{d^2 Z}{dz^2} - k^2 Z = 0 \quad (4.4)$$

$$\text{so } Z = A \exp(kz) + B \exp(-kz)$$

$$\text{and } \frac{1}{R} \frac{d}{dR} \left(R \frac{dJ}{dR} \right) + k^2 J(R) = 0 \quad (4.5)$$

Potential due to thin disk

Bessel Functions

$$\frac{1}{R} \frac{d}{dR} \left(R \frac{dJ}{dR} \right) + k^2 J(R) = 0 \quad 4.5$$

We would quite like $\Phi(R, \infty)$ and $\Phi(R, -\infty)$ to be zero, so

$$Z(z) = A \exp(-k|z|)$$

is the appropriate solution for $Z(z)$.

The R equation (4.5) is the defining equation for a Bessel function. These are the analogues of sines and cosines now for cylindrical as opposed to linear problems (e.g. drum beats).

So while $\frac{d^2 y}{dz^2} + k^2 y = 0$ has solutions $\sin(kz)$, $\cos(kz)$

$$(4.6)$$

similarly $\frac{1}{s} \frac{d}{ds} \left(s \frac{dy}{ds} \right) + k^2 y = 0$ has solutions $J_0(ks)$, $Y_0(ks)$

$$(4.7)$$

see e.g. Abramowitz & Stegun "Handbook of Mathematical Functions"

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk

Bessel Functions

Note that as $x \rightarrow 0$ $J_0(x) \rightarrow 1$ and $Y_0(x) \rightarrow -\infty$.

More generally the equation

$$\frac{1}{s} \frac{d}{ds} \left(s \frac{dy}{ds} \right) + \left(k^2 - \frac{\nu^2}{s^2} \right) y = 0$$

has solutions $J_\nu(ks)$, $Y_\nu(ks)$, a family of Bessel functions characterized by the index ν .

J_ν - Bessel function of the first kind

Y_ν - Bessel function of the second kind

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy


Spiral Galaxy Rotation Curves

Potential due to thin disk

Bessel Functions



J_ν , Bessel function of the first kind



Y_ν , Bessel function of the second kind

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk

Bessel Functions

$$\text{Similarly} \quad \frac{1}{s} \frac{d}{ds} \left(s \frac{dy}{ds} \right) - k^2 y = 0 \rightarrow I_0(ks), K_0(ks)$$

$$\text{and} \quad \frac{1}{s} \frac{d}{ds} \left(s \frac{dy}{ds} \right) - \left(k^2 + \frac{\nu^2}{s^2} \right) y = 0 \rightarrow I_\nu(ks), K_\nu(ks)$$

see Abramowitz + Stegun "Handbook of Mathematical functions"

These are **Modified Bessel Functions**.

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy


Spiral Galaxy Rotation Curves

Potential due to thin disk

Modified Bessel Functions



I_ν , Modified Bessel function of the first kind



K_ν , Modified Bessel function of the second kind

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk

Bessel Functions

And we can take this even further. By analogy with Fourier transforms where $\sin, \cos \rightarrow$ form the basis, we have $J, Y \rightarrow$ Hankel transforms.

Given a function $g(r)$, then the Hankel transform of g is

$$\tilde{g}(k) = \int_0^{\infty} g(r) J_{\nu}(kr) r dr$$

and the inverse transform is:

$$g(r) = \int_0^{\infty} \tilde{g}(k) J_{\nu}(kr) k dk$$

look these up in books of Hankel transforms!

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



Potential due to thin disk

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

$$\frac{d^2 Z}{dz^2} - k^2 Z = 0 \quad 4.4$$

$$\frac{1}{R} \frac{d}{dR} \left(R \frac{dJ}{dR} \right) + k^2 J(R) = 0 \quad 4.5$$

Returning to the axisymmetric plane distribution, we have

$$(4.4) \Rightarrow Z(z) = \exp(-k|z|)$$

$$(4.5) \Rightarrow J(R) = J_0(kr)$$

choose J to get Φ finite at $R = 0$

Let $k > 0$ then

$$\Rightarrow \Phi_k(R, Z) = Ce^{-kz} J_0(kR) \quad z > 0$$

$$Ce^{kz} J_0(kR) \quad z < 0$$

This is true $\forall k > 0$, but a specific k for each Φ_k .

$$\text{General potential} \rightarrow \sum_k \Phi_k$$

Potential due to thin disk

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

$$\Rightarrow \Phi(R, z) = \int_0^{\infty} f(k) e^{-k|z|} J_0(kR) dk \quad (4.8)$$

Here $f(k)$ is a weighting function, corresponding to the C values in the sum. So what we need to do for a particular mass distribution is find $f(k)$.

If we are going to relate it to a mass distribution, the next thing we should do is look at the $z = 0$ plane, i.e. the region we have neglected so far since we have taken $\nabla^2 \Phi = 0$ and so considered regions outside the plane.

Note that Φ_k is continuous across $z = 0$ but $\nabla \Phi_k$ **is not**, due to $|z|$ dependence. That is where the mass is, so that is not a surprise.

$\Rightarrow \nabla^2 \Phi_k = 0$ except at $z = 0$ and $\Phi_k \rightarrow 0$ as $z, R \rightarrow \infty \Rightarrow$ satisfies conditions for potential from an isolated mass distribution. Still need to link with ρ (or $\Sigma(R)$) in the plane.

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk

Poisson's equation



Use Gauss' Theorem (\equiv Poisson's equation plus divergence theorem) to determine Σ in the $z = 0$ plane.

Over the cylinder

$$\begin{aligned} \iiint 4\pi G\rho dV &= \iiint \nabla^2\Phi dV = \iiint \nabla \cdot (\nabla\Phi) dV \\ &= \iint \nabla\Phi \cdot \hat{n} d^2\mathbf{S} \end{aligned}$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk

Poisson's equation

$$\iiint 4\pi G\rho dV = \iint \nabla\Phi \cdot \hat{n} d^2S$$

Consider the limit in which the cylinder height $\rightarrow 0$. Then if A is the area of an end of the cylinder

$$LHS = 4\pi G\Sigma A$$

$$RHS = \left(\left[\frac{\partial\Phi}{\partial z} \right]_{z=0+} - \left[\frac{\partial\Phi}{\partial z} \right]_{z=0-} \right) \times A$$

$$\text{Equating these} \Rightarrow \left[\frac{\partial\Phi}{\partial z} \right]_{0-}^{0+} = 4\pi G\Sigma(R)$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk

Poisson's equation

$$\begin{aligned}
 RHS &= - \int_0^{\infty} kf(k)e^{-k0+} J_0(kR)dk - \int_0^{\infty} kf(k)e^{-k0-} J_0(kR)dk \\
 &= - \int_0^{\infty} kf(k)J_0(kR)dk - \int_0^{\infty} kf(k)J_0(kR)dk \\
 &= -2 \int_0^{\infty} kf(k)J_0(kR)dk \\
 \Rightarrow \Sigma(R) &= -\frac{1}{2\pi G} \int_0^{\infty} f(k)J_0(kR)kdk
 \end{aligned}$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk

Poisson's equation

Hence determine $f(k)$ [and hence Φ] from inverse Hankel transform

$$f(k) = -2\pi G \int_0^\infty \Sigma(R) J_0(kR) R dR$$

Thus the process for determining Φ from ρ in this case is $\Sigma \rightarrow f \rightarrow \Phi$.

Note: For determining the circular velocity need $\frac{\partial \Phi}{\partial R}$, which becomes $\frac{dJ_0(x)}{dx}$, and for Bessel function J_0 have $\frac{d}{dx} J_0(x) = -J_1(x)$ [Example].

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk

Summary of derivation of Φ for thin axisymmetric disk

This has been a bit longwinded, but the steps are clear. They are:

- ① $\nabla^2\Phi = 0$ outside disk. Solve by separation of variables.
- ② Solutions of form $\Phi_k(R, z) = Ce^{-k|z|}J_0(kR) \forall k > 0$
- ③ $\Phi_k \rightarrow 0$ as $R, z \rightarrow \infty$ **and** satisfies $\nabla^2\Phi = 0$
 \Rightarrow is potential of an isolated density distribution
- ④ General Φ can be written as

$$\Phi(R, z) = \int_0^\infty \Phi_k(R, z)f(k)dk$$

where $f(k)$ is an appropriate weight function.

- ⑤ Use Gauss' theorem to determine

$$\Sigma(R) = -\frac{1}{2\pi G} \int_0^\infty f(k)J_0(kR)kdk$$

- ⑥ Hence

$$f(k) = -2\pi G \int_0^\infty \Sigma(R)J_0(kR)RdR$$

So given Σ , use item (6) to determine $f(k)$, and then (5) to obtain Φ .

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions
Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potential due to thin disk

Circular velocity in the thin disk

$$\Phi(R, z) = \int_0^\infty f(k) e^{-k|z|} J_0(kR) dk \quad 4.8$$

The circular velocity in the plane of a plane distribution of matter is given by

$$\frac{v_C^2(R)}{R} = \left. \frac{\partial \Phi}{\partial R} \right|_{z=0}$$

and we have

$$x = kR \quad \frac{d}{dR} J_0(kR) = k \frac{d}{dx} J_0 x = -kJ_1(kR)$$

$$\frac{d}{dx} [x^m J_m(x)] = x^m J_{m-1}(x)$$

Then since we have equation (4.8), then

$$\frac{v_C^2(R)}{R} = - \int_0^\infty f(k) J_1(kR) k dk$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy


Spiral Galaxy Rotation Curves

Thin Disk: Examples

Mestel disk

A Mestel disk has the surface density distribution

$$\Sigma(R) = \frac{\Sigma_0 R_0}{R}$$



Thus

$$M(< R) = \int_0^R 2\pi \Sigma(R') R' dR' = 2\pi R_0 \Sigma_0 \int_0^R dR'$$

$$= 2\pi \Sigma_0 R_0 R$$

$$\rightarrow \infty \text{ as } R \rightarrow \infty$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions
Poisson's equation
Summary of derivation of Φ for thin axisymmetric disk
Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Thin Disk: Examples

Mestel disk

$$f(k) = -2\pi G \int_0^\infty \Sigma(R) J_0(kR) R dR$$

$$\begin{aligned} f(k) &= -2\pi G \Sigma_0 R_0 \int_0^\infty J_0(kR) dR \\ &= -\frac{2\pi G \Sigma_0 R_0}{k} \end{aligned}$$

$$\text{From Gradshteyn and Ryzhik 6.511.1 } \int_0^\infty J_\nu(bx) dx = \frac{1}{b} \quad \begin{matrix} \text{Re}(\nu) > -1 \\ b > 0 \end{matrix}$$

$$\Rightarrow \Phi(R, z) = -2\pi G \Sigma_0 R_0 \int_0^\infty e^{-k|z|} \frac{J_0(kR)}{k} dk$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions
Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Thin Disk: Examples

Mestel disk

$$\frac{v_c^2(R)}{R} = - \int_0^\infty f(k) J_1(kR) k dk$$

$$f(k) = - \frac{2\pi G \Sigma_0 R_0}{k}$$

$$\text{and } \frac{v_c^2(R)}{R} = 2\pi G \Sigma_0 R_0 \int_0^\infty J_1(kR) dk$$

$$\Rightarrow v_c^2(R) = 2\pi G \Sigma_0 R_0 = \text{const}$$

$$M(< R) = 2\pi \Sigma_0 R_0 R$$

Note that

$$v_c^2(R) = \frac{GM(R)}{R}$$

exactly in this case even though distribution is a disk, not spherical. More generally, find $v_c^2 \equiv \frac{GM(R)}{R}$ to within 10% [reasonable accuracy] for most smooth Σ distributions.

Conclude that measurement of $v_c(R)$ is a good measure of mass **inside** R .

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions
Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples
Mestel disk
Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Thin Disk: Examples

Exponential Disk

Here

$$\Sigma(R) = \Sigma_0 \exp[-R/R_d]$$

This has finite mass

$$\begin{aligned} M &= \int_0^{\infty} 2\pi \Sigma_0 \exp[-R/R_d] R dR \\ &= 2\pi \Sigma_0 R_d^2 \underbrace{\int_0^{\infty} e^{-x} x dx}_{=1} \\ &= 2\pi \Sigma_0 R_d^2 \end{aligned}$$

Then

$$f(k) = -2\pi G \Sigma_0 \int_0^{\infty} e^{-R/R_d} J_0(kR) R dR$$

$$\text{Gradshteyn + Ryzhik } \int_0^{\infty} e^{-\alpha x} J_0(\beta x) x dx = \frac{\alpha}{[\beta^2 + \alpha^2]^{3/2}}$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions
Poisson's equation
Summary of derivation of Φ for thin axisymmetric disk
Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Thin Disk: Examples

Exponential Disk

[Actually they have something (6.623.2) which requires a little work:

$$\int_0^{\infty} e^{-\alpha x} J_{\nu}(\beta x) x^{\nu+1} dx = \frac{2\alpha(2\beta)^{\nu} \Gamma(\nu + \frac{3}{2})}{\sqrt{\pi} (\alpha^2 + \beta^2)^{\nu + \frac{3}{2}}}$$

and you need to put in $\nu = 0$, $\Gamma(3/2) = \sqrt{\pi}/2$.

Then put $\alpha = 1/R_d$ and $\beta = k$.]

$$\int_0^{\infty} e^{-\alpha x} J_0(\beta x) x dx = \frac{\alpha}{[\beta^2 + \alpha^2]^{3/2}}$$

OK, so

$$f(k) = -\frac{2\pi G \Sigma_0 R_d^2}{[1 + (kR_d)^2]^{3/2}}$$

Hence

$$\Phi(R, z) = -2\pi G \Sigma_0 R_d^2 \int_0^{\infty} \frac{J_0(kR) e^{-k|z|}}{(1 + (kR_d)^2)^{3/2}} dk$$

Thin Disk: Examples

Exponential Disk

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions
Poisson's equation
Summary of derivation of Φ for thin axisymmetric disk
Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Use

$$\int_0^{\infty} \frac{J_{\nu}(xy) dx}{(x^2 + a^2)^{1/2}} = I_{\nu/2} \left(\frac{1}{2} ay \right) K_{\nu/2} \left(\frac{1}{2} ay \right)$$

You can do this with help from Gradshteyn + Ryzhik again, using (6.552.1)

$\frac{d}{da}$ of this gives

$$-a \int_0^{\infty} \frac{J_{\nu}(xy) dx}{(x^2 + a^2)^{3/2}} = \frac{y}{2} I_{\nu/2} \left(\frac{1}{2} ay \right) K'_{\nu/2} \left(\frac{1}{2} ay \right) + \frac{y}{2} I'_{\nu/2} \left(\frac{1}{2} ay \right) K_{\nu/2} \left(\frac{1}{2} ay \right)$$

so for $\nu = 0$

$$-a \int_0^{\infty} \frac{J_0(xy) dx}{(x^2 + a^2)^{3/2}} = -\frac{y}{2} I_0 \left(\frac{1}{2} ay \right) K_1 \left(\frac{1}{2} ay \right) + \frac{y}{2} I_1 \left(\frac{1}{2} ay \right) K_0 \left(\frac{1}{2} ay \right)$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Thin Disk: Examples

Exponential Disk

or

$$\int_0^\infty \frac{J_\nu(xy) dx}{(x^2 + a^2)^{3/2}} = \frac{y}{2a} \left[I_0\left(\frac{1}{2}ay\right)K_1\left(\frac{1}{2}ay\right) - I_1\left(\frac{1}{2}ay\right)K_0\left(\frac{1}{2}ay\right) \right]$$

Then with $x = k$, $y = R$ and $a = 1/R_d$ this becomes

$$\int_0^\infty \frac{J_\nu(kR) dk}{(1 + (kRd)^2)^{3/2}} = \frac{R}{2R_d^2} \left[I_0\left(\frac{R}{2R_d}\right)K_1\left(\frac{R}{2R_d}\right) - I_1\left(\frac{R}{2R_d}\right)K_0\left(\frac{R}{2R_d}\right) \right]$$

or

$$\Phi(R, 0) = -\pi G \Sigma_0 R \left[I_0\left(\frac{R}{2R_d}\right)K_1\left(\frac{R}{2R_d}\right) - I_1\left(\frac{R}{2R_d}\right)K_0\left(\frac{R}{2R_d}\right) \right]$$

Also you find for the circular velocity (with $y = R/2R_d$)

$$v_C^2 = R \frac{\partial \Phi}{\partial R} = 4\pi \Sigma_0 R_d y^2 [I_0 K_0 - I_1 K_1]$$

which is helpfully left as an example...

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

Thin Disk: Examples

Exponential disk




Figure 2.16 Contours in the (R, z) plane of constant potential Φ for a razor-thin exponential disk. The contour levels are GM_d/R_d divided by 1.5, 2, 2.5, ..., where $M_d = 2\pi\Sigma_0 R_d^2$ is the mass of the disk.

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions
Poisson's equation
Summary of derivation of Φ for thin axisymmetric disk

Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Thin Disk: Examples

Exponential disk




Figure 2.17 The circular-speed curves of: an exponential disk (full curve); a point mass with the same total mass (dotted curve); the spherical body for which $M(r)$ is given by equation (2.166) (dashed curve).

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions

Poisson's equation

Summary of derivation of Φ for thin axisymmetric disk

Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Thin Disk: Examples

Kuzmin disk




Figure 6.1: Surgery on the potential of a point mass produces the potential of a Kuzmin disk. Left: potential of a point mass. Contours show equal steps of $\Phi \propto 1/r$, while the arrows in the upper left quadrant show the radial force field. Dotted lines show $z = \pm a$. Right: potential of a Kuzmin disk, produced by excising the region $|z| \leq a$ from the field shown on the left. Arrows again indicate the force field; note that these no longer converge on the origin.

Image credit: Joshua Barnes

Potentials due to axisymmetric matter distributions

Potential due to thin disk

Bessel Functions
 Poisson's equation
 Summary of derivation of Φ for thin axisymmetric disk
 Examples

Mestel disk

Exponential Disk

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Thin Disk: Examples

Kuzmin disk



$$\Phi = \frac{-GM}{\sqrt{R^2 + (a + |z|)^2}}$$

Most (over)used disk model in astronomy is that credited to Miyamoto-Nagai (>1000 citations), which is a slight modification of Kuzmin disk.

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants: the rotation of our Galaxy

The Galaxy in the optical



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants: the rotation of our Galaxy

The Galaxy in the infrared



The Galactic coordinate system

The Oort Constants: the rotation of our Galaxy

Potentials due to axisymmetric matter distributions

Potential due to thin disk


The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Suppose all stars are in circular orbits, with velocities $v(R) = R\Omega(R)$
 Observe a star at distance d , galactic longitude ℓ .



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants: the rotation of our Galaxy

Line of sight velocity



The radial velocity, as seen from the Earth, is

$$v_R = v \cos \alpha - v_0 \sin \ell$$

But

$$\frac{\sin \ell}{R} = \frac{\sin \left(\frac{\pi}{2} + \alpha \right)}{R_0} = \frac{\cos \alpha}{R_0}$$

and so

$$v_R = \left(\frac{vR_0}{R} - v_0 \right) \sin \ell = \left(\frac{v}{R} - \frac{v_0}{R_0} \right) R_0 \sin \ell$$

Now if we look at only nearby stars ($d \ll R_0$):

$$R_0 - R \approx d \cos \ell$$

Potentials due to axisymmetric matter distributions


Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



$$v_R = \left(\frac{v}{R} - \frac{v_0}{R_0} \right) R_0 \sin \ell$$

$$R_0 - R \approx d \cos \ell$$

The Oort Constants: the rotation of our Galaxy

Line of sight velocity

If we expand $\frac{v}{R}$ around $R = R_0$:

$$\frac{v}{R} = \frac{v_0}{R_0} + (R - R_0) \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0}$$

So the radial velocity

$$v_R \approx -R_0 \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0} d \sin \ell \cos \ell$$

Since $\sin 2\ell = 2 \sin \ell \cos \ell$, we have

$$v_R = A d \sin 2\ell$$

$$A = -\frac{R_0}{2} \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0} = \frac{1}{2} \left[\frac{v_0}{R_0} - \frac{dv}{dR} \Big|_{R_0} \right]$$

the Oort constant A

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants: the rotation of our Galaxy

Line of sight velocity



$$v_R = Ad \sin 2\ell$$

$$A = -\frac{R_0}{2} \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0} = \frac{1}{2} \left[\frac{v_0}{R_0} - \frac{dv}{dR} \Big|_{R_0} \right]$$

Then if we measure $\frac{v_R}{d}$ as a function of ℓ , then (in principle) we would know A .

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants: the rotation of our Galaxy

Tangential velocity



Similarly we can measure the tangential velocity (from proper motions)

$$v_T = v \sin \alpha - v_0 \cos \ell$$

Note that $R \sin \alpha = R_0 \cos \ell - d$, and so

$$v_T = \frac{v}{R} [R_0 \cos \ell - d] - v_0 \cos \ell$$

i.e.

$$v_T = \left(\frac{v}{R} - \frac{v_0}{R_0} \right) R_0 \cos \ell - \frac{v}{R} d$$

For small d we have $R_0 - R \approx d \cos \ell$ and

$$\frac{v}{R} = \frac{v_0}{R_0} + (R - R_0) \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0} \quad (*)$$

The Oort Constants: the rotation of our Galaxy

Tangential velocity

Potentials due to axisymmetric matter distributions


Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



$$v_T = \left(\frac{v}{R} - \frac{v_0}{R_0} \right) R_0 \cos \ell - \frac{v}{R} d$$

$$\frac{v}{R} - \frac{v_0}{R_0} = (R - R_0) \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0} \quad (*)$$

$$R_0 - R \approx d \cos \ell$$

$$v_T \approx -R_0 \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0} d \cos^2 \ell - \frac{v}{R} d$$

The Oort Constants: the rotation of our Galaxy

Tangential velocity

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

$$\frac{v}{R} = \frac{v_0}{R_0} + (R - R_0) \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0} \quad (*)$$

We can remove the $\frac{v}{R}$ term by using the expression (*), and $\cos^2 \ell = \frac{1}{2}(1 + \cos 2\ell)$ to obtain

$$\begin{aligned} v_T &\approx -R_0 \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0} d \cos^2 \ell - \frac{v}{R} d \\ &\approx \frac{1}{2} \left(\frac{v_0}{R_0} - \frac{dv}{dR} \Big|_{R_0} \right) d \cos 2\ell - \frac{1}{2} \left(\frac{v_0}{R_0} + \frac{dv}{dR} \Big|_{R_0} \right) d + O\left(\frac{d^2}{R_0^2}\right) \end{aligned}$$

Now define

$$v_T = d(A \cos 2\ell + B)$$

i.e.

$$B = -\frac{1}{2} \left[\frac{v_0}{R_0} + \frac{dv}{dR} \Big|_{R_0} \right] \quad A = \frac{1}{2} \left[\frac{v_0}{R_0} - \frac{dv}{dR} \Big|_{R_0} \right]$$

The Oort Constants: the rotation of our Galaxy

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

$$A = \frac{1}{2} \left[\frac{v_0}{R_0} - \left. \frac{dv}{dR} \right|_{R_0} \right] = -\frac{R_0}{2} \frac{d}{dR} \left(\frac{v}{R} \right) \Big|_{R_0} = -\frac{1}{2} R_0 \left. \frac{d\Omega}{dR} \right|_{R_0}$$

$$B = -\frac{1}{2} \left[\frac{v_0}{R_0} + \left. \frac{dv}{dR} \right|_{R_0} \right] = -\left(\Omega + \frac{1}{2} R_0 \left. \frac{d\Omega}{dR} \right|_{R_0} \right)$$

- A measures the shear in the disk i.e. deviation from solid body rotation. For solid body rotation $\frac{d\Omega}{dR} = 0$, so $A = 0$.
- B measures vorticity i.e. tendency for material in the disk to circulate due to differential rotation.

Then note that

$$\Omega_0 = v_0/R_0 = A - B$$

and

$$\left. \frac{dv}{dR} \right|_{R_0} = -(A + B)$$

The Oort Constants: the rotation of our Galaxy

Measurement

Potentials due to axisymmetric matter distributions


Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



The Oort Constants: the rotation of our Galaxy

Measurement

Potentials due to axisymmetric matter distributions


Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



Not easy: distances, peculiar velocities, proper motions

The Oort Constants: the rotation of our Galaxy

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Best estimates are $A = 14.5 \text{ km s}^{-1} \text{ kpc}^{-1}$, $B = -12 \text{ km s}^{-1} \text{ kpc}^{-1}$.

$\Rightarrow \Omega_{R_0} = 26 \text{ km s}^{-1} \text{ kpc}^{-1}$, so with $R_0 = 8.5 \text{ kpc}$ $v_0 = 220 \text{ km s}^{-1}$

\Rightarrow Rotation period = 2.4×10^8 years.

Epicyclic frequency $K_0 = \sqrt{-4B(A - B)} = 36 \pm 10 \text{ km s}^{-1} \text{ kpc}^{-1}$

$K_0/\Omega_0 = 1.3 \pm 0.2$.

$$K^2 = R \frac{d\Omega^2}{dR} + 4\Omega^2$$

See examples sheet 3

The Sun makes 1.3 oscillations in the radial direction in the time it takes to complete the orbit around the Galactic centre

The Oort Constants: the rotation of our Galaxy

$$\nu^2 = \left(\frac{\partial^2 \Phi}{\partial z^2} \right)$$

$$\Omega^2 = \frac{1}{R} \left(\frac{\partial \Phi}{\partial R} \right)$$

Poisson's equation in cylindrical coordinates:

$$4\pi G\rho = \frac{1}{R} \frac{\partial}{\partial R} \left(R \frac{\partial \Phi}{\partial R} \right) + \frac{\partial^2 \Phi}{\partial z^2}$$

$$\approx \frac{1}{R} \frac{dv_c^2}{dR} + \nu^2$$

If we could measure ν , the vertical frequency, we could estimate the mass density in the Galactic plane, but we can't.

But, if we assume that the matter distribution was spherical then mass internal to the Sun:

$$\frac{v_0^2}{R_0} \approx \frac{GM}{R_0^2}$$

$$\Rightarrow M \approx \frac{R_0 v_0^2}{G} \approx 10^{11} M_\odot$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants: the rotation of our Galaxy

Jan Oort, 1930s



Potentials due to axisymmetric matter distributions

Potential due to thin disk


The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants and epicycles



A star on an orbit with the guiding center radius R_g :

$$x \equiv R - R_g$$

The angular speed of the circular orbit

$$\Omega_g = L_z / R_g^2$$

$$\begin{aligned} \dot{\phi} &= \frac{p_\phi}{R^2} = \frac{L_z}{R_g^2} \left(1 + \frac{x}{R_g}\right)^{-2} \\ &\simeq \Omega_g \left(1 - \frac{2x}{R_g}\right) \end{aligned}$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants and epicycles

$$\dot{\phi} \simeq \Omega_g \left(1 - \frac{2x}{R_g} \right)$$

If we could measure $v_\phi(R_0) - v_c(R_0)$ and v_R for a group of stars, each with its own guiding center, is there a way to obtain the relevant frequencies?

$$\begin{aligned} v_\phi(R_0) - v_c(R_0) &= R_0(\dot{\phi} - \Omega_0) = R_0(\dot{\phi} - \Omega_g + \Omega_g - \Omega_0) \\ &\simeq R_0 \left[(\dot{\phi} - \Omega_g) - \left(\frac{d\Omega}{dR} \right)_{R_g} x \right] \end{aligned}$$

$$v_\phi(R_0) - v_c(R_0) \simeq -R_0 x \left(\frac{2\Omega}{R} + \frac{d\Omega}{dR} \right)_{R_g}$$

Evaluate at R_0 and dropping terms of order of x^2

$$v_\phi(R_0) - v_c(R_0) \simeq -x \left(2\Omega + R \frac{d\Omega}{dR} \right)_{R_0}$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants and epicycles

The Oort's constants definition:

$$A(R) \equiv \frac{1}{2} \left(\frac{v_c}{R} - \frac{dv_c}{dR} \right) = -\frac{1}{2} R \frac{d\Omega}{dR}$$

$$B(R) \equiv -\frac{1}{2} \left(\frac{v_c}{R} + \frac{dv_c}{dR} \right) = -\left(\Omega + \frac{1}{2} R \frac{d\Omega}{dR} \right)$$

Therefore,

$$v_\phi(R_0) - v_c(R_0) \simeq -x \left(2\Omega + R \frac{d\Omega}{dR} \right)_{R_0} = 2Bx$$

Epicyclic radial motion solution:

$$x(t) = X \cos(Kt + \alpha)$$

Hence, averaging over the phases of stars near the Sun

$$\overline{[v_\phi - v_c(R_0)]^2} = 2B^2 X^2$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Oort Constants and epicycles

$$\overline{[v_\phi - v_c(R_0)]^2} = 2B^2 X^2$$

Similarly, for the radial velocity

$$\overline{v_R^2} = -2B(A - B)X^2$$

Taking ratio:

$$\frac{\overline{[v_\phi - v_c(R_0)]^2}}{\overline{v_R^2}} \simeq \frac{-B}{A - B} = -\frac{B}{\Omega_0} = \frac{K_0^2}{4\Omega_0^2}$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Rotation Curve of our Galaxy

21cm line of neutral hydrogen



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Addison-Wesley.

frequency 1420.4058 MHz, wavelength 21 cm, probability **1 in 10^7 years**

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Rotation Curve of our Galaxy

21cm line of neutral hydrogen



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Rotation Curve of our Galaxy


21cm line of neutral hydrogen

TIDAL INTERACTIONS IN M81 GROUP

Stellar Light Distribution



21 cm HI Distribution



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Rotation Curve of our Galaxy

Rotation curve from 21cm line




Figure 9.2 The traces in the (l, v) plot of three rings. The radii of the rings are $r = 0.9R_0$ (full curve), $r = 0.1R_0$ (dotted curve) and $r = 1.5R_0$ (dashed curve). The circular speed in the disk is unity at all R .

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Rotation Curve of our Galaxy

Rotation curve from 21cm line




Figure 9.3 Grayscale of optical depth in the (l, v) plane. As for Figure 8.12, the number density of hydrogen atoms has been assumed to equal $2 \times 10^5 \text{ m}^{-3}$ at R_0 and to vary with R as $\exp(-R/2R_0)$. The circular speed is given by $v_c(R) = (R/R_0)^{0.1} \times 210 \text{ km s}^{-1}$. The sine curve $v_{\text{los}} = -v_c(R_0) \sin l$ (shown dashed) lies just outside the boundary of the occupied band. Figure 9.4 shows a section of the surface $\tau(l, v)$ along the vertical dotted line $l = 20^\circ$.

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Rotation Curve of our Galaxy

Terminal velocity

Gas moves on circular orbits - use 21cm HI emission, and measure Doppler shift given ℓ .

We have

$$v_R = \left(\frac{v}{R} - \frac{v_0}{R_0} \right) R_0 \sin \ell$$

If $\Omega(R) = \frac{v}{R}$ decreases monotonically with R , then we can look at the 21cm line profile and pick the maximum Doppler shift as the one corresponding to the smallest circle along the line of sight at Galactic longitude ℓ . This occurs when $R = R_0 \sin \ell$

Therefore

$$v_R(\max) = v(R) - v_0 \sin \ell$$

and hence

$$v(R) = v_R(\max) + v_0 \sin \ell$$

Note that this works only for $-90^\circ < \ell < 90^\circ$, since a line of sight for $90^\circ < \ell < 270^\circ$ is never a tangent to a circular orbit.

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Rotation Curve of our Galaxy

Non-circular motion

- Axisymmetric expansion
- Oval distortion (elliptical orbits)
- Spiral structure
- Random motions

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Rotation Curve of our Galaxy

Axisymmetric expansion




Figure 9.6 The (l, v) trace of a ring interior to the solar circle that is expanding radially as well as rotating. The expansion velocity is just $1/20$ of the circular speed in the disk. The full curve in Figure 9.2 shows the trace of the ring in the absence of expansion.

Potentials due to
axisymmetric matter
distributions

Potential due to thin disk

The Oort Constants: the
rotation of our Galaxy





The Oort Constants and
epicycles

The Rotation Curve of
our Galaxy

Spiral Galaxy Rotation
Curves

The Rotation Curve of our Galaxy

Oval distortion and spiral structure



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy


The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

The Rotation Curve of our Galaxy

Rotation curve from 21cm line



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

- spirals contain disks of gas. In a symmetric potential settles down to rotate on circular orbits in centrifugal balance
- gas collides with itself and shocks, therefore paths of gas motion rarely intersect, unlike stars
- if the potential is spherically symmetric then the measure of the circular velocity $v_c(R)$ is a good measure of the mass $M(R)$ contained within radius R

$$\frac{v_c^2}{R} = \frac{d\Phi}{dR} \simeq \frac{GM(R)}{R^2}$$

or

$$M(< R) \simeq \frac{v_c^2 R}{G}$$

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

NGC 253



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

We can obtain $v_C(R)$ by observing the Doppler shift of a spectral line. For the gas in the spiral's disk, our options are:

- HI - **radio** - neutral gas, largest range in radius
- H_α - **optical** - warm gas, inner regions
- CO - **mm** - molecular gas, very inner regions

Problems:

- **Instrumental *beam smearing***: each pointing has data from a range of radii, need to deconvolve
- **Intrinsic**
 - absorption*: especially if galaxy is close to edge on position
 - finite thickness of gas layer*

Spiral arms cause kinks in apparent rotation curve, because

- gas motion is not exactly circular
- arms provide non-axisymmetric potential
- arms contribute to optical extinction

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

NGC 253 rotation curve




FIG. 2a-1




FIG. 2a-2




FIG. 2b

FIG. 2.—(a-1) Inner rotation curve of NGC 253 derived by using the CO PV diagram, and (a-2) CO + optical (Hz) (= total) rotation curve compared to H I rotation. (b) Total rotation curve of NGC 253 obtained by averaging and smoothing the eastern and western halves of the rotation velocities. See Table 1 for observational parameters and references.

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

NGC 3079



Spiral Galaxy Rotation Curves

NGC 3079 rotation curve from PV diagram

Potentials due to axisymmetric matter distributions



Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

NGC 4565



Spiral Galaxy Rotation Curves

NGC 6946

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

NGC 1808



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

Rotation curves of nearby spirals




FIG. 12.—Rotation curves of galaxies studied in this paper plotted in the same linear and velocity scales. (a) Rotation curves having a central peak similar to that of our Galaxy. (b) Rotation curves without significant central peak.

Spiral Galaxy Rotation Curves

IC 342

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

NGC 5907



Spiral Galaxy Rotation Curves

NGC 5907's secret

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves



Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

Vera Rubin, 1980



“...This form for the rotation curves implies that the mass is not centrally condensed, but that significant mass is located at large R . The integral mass is increasing at least as fast as R . The mass is not converging to a limiting mass at the edge of the optical image. The conclusion is inescapable that non-luminous matter exists beyond the optical galaxy...”

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

The similarity of the rotation curves




Figure 4 Rotation curves of spiral galaxies obtained by combining CO data for the central regions, optical for disks, and HI for outer disk and halo (Sofue et al. 1999a).

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

The dynamical components




Fig. 4. Least-squares fit by the bulge, disk, and dark halo to the grand rotation curve. The thick line represents the fitted rotation curve, and the thin lines show individual contributions from bulge, disk, and halo. The observed velocities, $V(R_i)$, are shown by open circles, and the most recent accurate results from VERA (Honma et al. 2007; Oh et al. 2010) are shown by squares.

Potentials due to axisymmetric matter distributions

Potential due to thin disk

The Oort Constants: the rotation of our Galaxy

The Oort Constants and epicycles

The Rotation Curve of our Galaxy

Spiral Galaxy Rotation Curves

Spiral Galaxy Rotation Curves

Tully-Fisher relation

